

APCI 2022-2023 Flu Vaccine Pre-Book Order

MAKE YOUR PRE-BOOK RESERVATION BY **MAY 20, 2022**
FOR THE 2022-2023 FLU SEASON TO RECEIVE THE BELOW OFFER.*



FIRM DOSE DISCOUNT

3% discount
on orders placed by
April 15, 2022



PROMPT PAY DISCOUNT

2% discount
on orders paid
within 60 days



RISK SHARING

10% return of each brand
ordered and unused by the
end of the season

2022-2023 INFLUENZA VACCINE
DELIVERY EXPECTATIONS:

9/2/2022


35%

9/30/2022

75%

10/15/2022

100%

	 QUADRIVALENT influenza vaccine, adjuvanted	 Influenza Vaccine FLUCELVAX QUADRIVALENT	 INFLUENZA VACCINE afluria QUADRIVALENT	
	65+ years	6+ months		6+ months
	0.5-mL Pre-filled Syringe	0.5-mL Pre-filled Syringe	5-mL Multi-dose Vial	5-mL Multi-dose Vial
				36+ months
				0.5-mL Pre-filled Syringe
NET PRICE/UNIT*	\$487.92	\$200.93	\$192.17	\$154.46
				\$165.07

*Inclusive of firm dose discount. Does not include federal excise tax. Additional discounts may apply

When it comes to pre-booking, we make it easy—because **we've got flu covered™**.

Visit flu360.com to place your pre-book reservation, manage your account, and access vaccination resources, complimentary to you as a partner of Seqirus.

**FLUAD® QUADRIVALENT
(Influenza Vaccine, Adjuvanted)
INDICATION and IMPORTANT SAFETY
INFORMATION**

INDICATION AND USAGE

FLUAD® QUADRIVALENT is an inactivated influenza vaccine indicated for active immunization against influenza disease caused by influenza virus subtypes A and types B contained in the vaccine. FLUAD QUADRIVALENT is approved for use in persons 65 years of age and older. This indication is approved under accelerated approval based on the immune response elicited by FLUAD QUADRIVALENT.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

Do not administer FLUAD QUADRIVALENT to anyone with a history of severe allergic reaction (e.g. anaphylaxis) to any component of the vaccine, including egg protein, or to a previous influenza vaccine.

WARNINGS AND PRECAUTIONS

If Guillain-Barré syndrome (GBS) has occurred within 6 weeks of receipt of prior influenza vaccine, the decision to give FLUAD QUADRIVALENT should be based on careful consideration of the potential benefits and risks.

Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of the vaccine.

The immune response to FLUAD QUADRIVALENT in immunocompromised persons, including individuals receiving immunosuppressive therapy, may be lower than in immunocompetent individuals.

Syncope (fainting) may occur in association with administration of injectable vaccines including FLUAD QUADRIVALENT. Ensure procedures are in place to avoid injury from falling associated with syncope.

Vaccination with FLUAD QUADRIVALENT may not protect all vaccine recipients against influenza disease.

ADVERSE REACTIONS

The most common (≥ 10%) local and systemic reactions in elderly subjects 65 years of age and older were injection site pain (16.3%), headache (10.8%) and fatigue (10.5%).

To report SUSPECTED ADVERSE REACTIONS, contact Seqirus USA Inc. at 1-855-358-8966 or VAERS at 1-800-822-7967 or www.vaers.hhs.gov.

Before administration, please see the [full US Prescribing Information](#) for FLUAD QUADRIVALENT.

FLUAD® QUADRIVALENT is a registered trademark of Seqirus UK Limited or its affiliates.

**FLUCELVAX® QUADRIVALENT (Influenza Vaccine)
IMPORTANT SAFETY INFORMATION**

INDICATION AND USAGE

FLUCELVAX QUADRIVALENT is an inactivated vaccine indicated for active immunization for the prevention of influenza disease caused by influenza virus subtypes A and types B contained in the vaccine. FLUCELVAX QUADRIVALENT is approved for use in persons 6 months of age and older

CONTRAINDICATIONS

Do not administer FLUCELVAX QUADRIVALENT to anyone with a history of severe allergic reactions (e.g. anaphylaxis) to any component of the vaccine.

WARNINGS AND PRECAUTIONS

If Guillain-Barré syndrome has occurred within 6 weeks of receipt of a prior influenza vaccine, the decision to give FLUCELVAX QUADRIVALENT should be based on careful consideration of the potential benefits and risks.

Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of the vaccine.

Syncope (fainting) can occur in association with administration of injectable vaccines, including FLUCELVAX QUADRIVALENT. Syncope can be accompanied by transient neurological signs such as visual disturbance, paresthesia, and tonic-clonic limb movements. Procedures should be in place to avoid falling injury and to restore cerebral perfusion following syncope by maintaining a supine or Trendelenburg position.

After vaccination with FLUCELVAX QUADRIVALENT, immunocompromised individuals, including those receiving immunosuppressive therapy, may have a reduced immune response.

Vaccination with FLUCELVAX QUADRIVALENT may not protect all vaccine recipients against influenza disease.

ADVERSE REACTIONS

In children 6 months through 3 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were tenderness (27.9%), erythema (25.8%), induration (17.3%) and ecchymosis (10.7%). The most common systemic adverse reactions were irritability (27.9%), sleepiness (26.9%), diarrhea (17.9%) and change of eating habits (17.4%).

In children 2 through 8 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were tenderness (28.7%), pain (27.9%) and erythema (21.3%), induration (14.9%) and ecchymosis (10.0%). The most common systemic adverse reactions were sleepiness (14.9%), headache (13.8%), fatigue (13.8%), irritability (13.8%) and loss of appetite (10.6%).

In children and adolescents 9 through 17 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were injection site pain (21.7%), erythema (17.2%) and induration (10.5%). The most common systemic adverse reactions were headache (18.1%) and fatigue (17.0%).

In adults 18 through 64 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were pain (45.4%), erythema (13.4%) and induration (11.6%). The most common systemic adverse reactions were headache (18.7%), fatigue (17.8%) and myalgia (15.4%).

In adults ≥65 years of age who received FLUCELVAX QUADRIVALENT, the most commonly reported injection-site adverse reactions were pain (21.6%) and erythema (11.9%).

To report SUSPECTED ADVERSE REACTIONS, contact Seqirus at 1-855-358-8966 or VAERS at 1-800-822-7967 or www.vaers.hhs.gov.

Before administration, please see the [full US Prescribing Information](#) for FLUCELVAX QUADRIVALENT.

FLUCELVAX® QUADRIVALENT is a registered trademark of Seqirus UK Limited or its affiliates.

**AFLURIA® QUADRIVALENT (Influenza Vaccine)
INDICATION and IMPORTANT SAFETY
INFORMATION**

INDICATION AND USAGE

AFLURIA QUADRIVALENT is an inactivated influenza vaccine indicated for active immunization against influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.

AFLURIA QUADRIVALENT is approved for use in persons 6 months of age and older.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

AFLURIA QUADRIVALENT is contraindicated in individuals with known severe allergic reactions (e.g., anaphylaxis) to any component of the vaccine including egg protein, or to a previous dose of any influenza vaccine.

WARNINGS AND PRECAUTIONS

If Guillain-Barré Syndrome (GBS) has occurred within 6 weeks of previous influenza vaccination, the decision to give AFLURIA QUADRIVALENT should be based on careful consideration of the potential benefits and risks.

Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of the vaccine.

If AFLURIA QUADRIVALENT is administered to immunocompromised persons, including those receiving immunosuppressive therapy, the immune response may be diminished.

Vaccination with AFLURIA QUADRIVALENT may not protect all individuals.

ADVERSE REACTIONS

AFLURIA QUADRIVALENT administered by needle and syringe:

In adults 18 through 64 years, the most commonly reported injection-site adverse reaction was pain (≥ 40%). The most common systemic adverse events were myalgia and headache (≥ 20%).

In adults 65 years of age and older, the most commonly reported injection-site adverse reaction was pain (≥ 20%). The most common systemic adverse event was myalgia (≥ 10%).

In children 5 through 8 years, the most commonly reported injection-site adverse reactions were pain (≥ 50%), redness and swelling (≥ 10%). The most common systemic adverse event was headache (≥ 10%).

In children 9 through 17 years, the most commonly reported injection-site adverse reactions were pain (≥ 50%), redness and swelling (≥ 10%). The most common systemic adverse events were headache, myalgia, and malaise and fatigue (≥ 10%).

In children 6 months through 35 months of age, the most commonly reported injection-site reactions were pain and redness (≥ 20%). The most common systemic adverse events were irritability (≥ 30%), diarrhea and loss of appetite (≥ 20%).

In children 36 through 59 months of age, the most commonly reported injection site reactions were pain (≥ 30%) and redness (≥ 20%). The most commonly reported systemic adverse events were malaise and fatigue, and diarrhea (≥ 10%).

The safety experience with AFLURIA (trivalent formulation) is relevant to AFLURIA QUADRIVALENT because both vaccines are manufactured using the same process and have overlapping compositions:

In adults 18 through 64 years of age, the most commonly reported injection-site adverse reactions with AFLURIA (trivalent formulation) when administered by the PharmaJet Stratis Needle-Free Injection System were tenderness (≥ 80%), swelling, pain, redness (≥ 60%), itching (≥ 20%) and bruising (≥ 10%). The most common systemic adverse events were myalgia, malaise (≥ 30%), and headache (≥ 20%).

To report SUSPECTED ADVERSE REACTIONS, contact Seqirus USA Inc. at 1-855-358-8966 or VAERS at 1-800-822-7967 or www.vaers.hhs.gov.

Before administration, please see the [full US Prescribing Information](#) for AFLURIA QUADRIVALENT.

AFLURIA® QUADRIVALENT is a registered trademark of Seqirus UK Limited or its affiliates. PharmaJet® and STRATIS® are registered trademarks of PharmaJet.